



Reproduction frequency, Growth forms, Growth Performance indices, Condition factors and Natural Mortality of Red Claw Crayfish (*Cherax quadricarinatus*, Von Martens, 1868) in the Kafue Floodplain Fishery, Zambia.

Hangoma Gordon Mudenda^a, Geoffrey Makwelele^b and Mauris Chinyama Makeche^{c,*}

^aDepartment of Biological Sciences, School of Natural Sciences, University of Zambia, Lusaka, Zambia.

^bDepartment of Biological Sciences, School of Mathematics and Science Education, Chalimbana University, Chongwe, Zambia.

^cInstitute of Basic and Biomedical Sciences, Levy Mwanawasa Medical University, Lusaka, Zambia.

ABSTRACT :

The research was conducted from the Kafue Floodplain Fishery because it is a World-recognised Ramsar-designated site that has diverse ecological habitats that are rich in fish biodiversity. Crayfish were sampled from Kafue Road Bridge and Chanyanya. A total of 462 *C. quadricarinatus* samples were collected from the Kafue Floodplain Fishery between November and December, 2022. Traps, locally called Kamono, were used to collect Crayfish. Length-weight relationships were determined by linear regression in Microsoft Excel to determine growth rates, r^2 , and allometric growth forms. Reproduction frequencies of the sampled Crayfish were determined using length-frequency histograms. The growth performance indices, Condition factors and natural mortality parameters were determined from the linear regression of the length-weight relationship. The results showed that Crayfish is a multiple spawner and it has a sex ratio of 2:1. The growth performance indices ranged between 4.3 to 4.9. All samples exhibited negative growth forms. Natural mortality coefficients were less than 1.0 and the Condition factors were all above 1, suggesting good adaptability to the habitat at Lake Kariba.

Keywords: *Cherax quadricarinatus*, reproduction, Condition factor, natural mortality.

Introduction

Cherax quadricarinatus, Red claw crayfish, is a freshwater crustacean belonging to the family Parastacidae (Austin *et al.*, 2010; FAO, 2011). Feral populations have been established in most tropical and subtropical freshwater systems across the globe from its native range of the tropical river catchments of Northern Australia and South-Eastern Papua New Guinea (Belle *et al.*, 2011). Aquaculture and ornamental aquarium trade are notably the main pathways behind the rapid spread of Crayfish globally. *C. quadricarinatus* is a hardy crustacean; it can be found in coastal streams and freshwater habitats, with a preference for the slower moving waters of rivers, lakes and lagoons (Austin *et al.*, 2010). *C. quadricarinatus* reproduction frequency is dependent on water temperature and day length, and normally spawns between September and April in freshwater bodies (Nunes *et al.*, 2016). *C. quadricarinatus* is a multiple spawner which is capable of spawning several times (3 to 5 times) each year as long as water temperatures remain above 18°C (Karplus *et al.*, 1997; Jaklic and Vrezec, 2011).

Length-weight studies are important in estimating various parameters on fish species such as reproduction frequency, growth forms, condition factors and natural mortalities (Osho and Usman, 2019). Length-weight relationships are also important in assessing the wellbeing of a fish populations (Makeche *et al.*, 2023). If the relationship between the length and weight of fish is isometric, it implies that the increase in fish weight is proportional to the cube of its length. Any other result depicts an allometric relationship, which can either be positive ($b > 3$) or negative allometric ($b < 3$) (Froese, 2006).

A Condition factor, which is obtained from length-weight measurements, can be used as an indicator of the health and sustainability of a fish stock. Various factors such as sex, seasons, environmental conditions and food availability all affect Condition factors of fish (Amponsah *et al.*, 2020). Condition factors are important in determining the ideal harvesting size, fish maturity and adaptability (Saha *et al.*, 2021). Fish species with Condition factor values greater than or equal to one (≥ 1) are in good condition and healthy while fish species with Condition factors less than one (< 1) are in a bad condition and unhealthy (Saha *et al.*, 2021).

Growth performance indices and growth rates are used in determining the type of growth a living organism exhibits. Growth of *C. quadricarinatus* is moderately fast ($r^2 > 0.5$) with longevity of approximately 4 years (Mukuka, 2019). This species has a growth performance index of between 2 (Makwelele, 2017) and 4 (Marufu *et al.*, 2018; Mukuka, 2019).

In Zambia, Crayfish are usually caught as a by catch (Makwelele, 2017). *C. quadricarinatus* death due to natural causes are low and usually result from diseases, competitors, spawning stress and other severe environmental factors (Nunes *et al.*, 2016). Makwelele (2017) reported natural mortality values

ranging from 0.06 to 0.08 from the Kafue Floodplains while Marufu *et al.* (2018) reported a natural mortality of 0.99 in Lake Kariba. Mukuka (2019) recorded an above-average natural mortality of 1.45 from the Upper Zambezi. Estimates of natural mortality parameters are important in improving the conservation of fish species (Hossain *et al.*, 2019; Saha *et al.*, 2021). Natural mortality is an important component which is used to estimate exploitation ratios of a fish stock in an aquatic habitat. An exploitation ratio which is less than 0.5 denotes under-exploitation while an exploitation ratio value above 0.5 depicts over-exploitation. A natural aquatic habitat is optimally exploited if its exploitation ratio is 0.5 (Makeche *et al.*, 2020).

The Kafue Floodplains are recognized as an internationally important Ramsar wetland due to the important biodiversity that this complex habitat supports (Ramsar, 2007). The Kafue Floodplains house two National parks (Lochinvar National Park and Blue Lagoon National Park) and it has abundant wild animals and various fish species such as Cichlids, Catfishes, *Hepsetus* spp, *Serranochromis* spp, *Labeo* spp, *Momyrus* and *Brycinus* spp among others (Cowx *et al.*, 2018). The Red claw crayfish was introduced in the Kafue Floodplains accidentally from fish farm escapees in the vicinity of Kafue Town in the 1990s (Tyser and Douthwaite, 2014).

The study had four main objectives: 1. to determine the reproduction frequency and sex ratio of *C. quadricarinatus* at Lake Kariba, 2. To determine the growth forms of *C. quadricarinatus* at Lake Kariba, 3. to determine the Condition factor of *C. quadricarinatus* at Lake Kariba, and 4. to determine the natural mortality coefficients of *C. quadricarinatus* at Lake Kariba.

Materials and Methods

2.1 Study sites and sampling

The research was conducted from the Kafue Floodplain Fishery because it is a World-recognized Ramsar site which is rich in biodiversity (Ramsar, 2007).

Crayfish were sampled from Chanyanya and Kafue Road Bridge (Fig. 1). Chanyanya lies along latitude $-15^{\circ}42'S$ and longitude $28^{\circ}00'E$. Kafue Road Bridge study site was located along latitude $-15^{\circ}84'S$ and longitude $28^{\circ}24'E$. A total of 462 *C. quadricarinatus* samples were collected from Lake Kariba between November and December, 2022.




Fig. 1 - The Kafue floodplain fishery with sampling stations (Source: modified From <http://www.bing.com/images>).

2.2 Data collection and analyses

Traps locally known as Kamono were used to trap sample Crayfish. The caught specimens were handled ethically by immediately immersing them in 10% formalin. The sampled Crayfish were weighed, measured and sexed. This sampling procedure was repeated for three nights at each study site. total length (TL), carapace length and Body weight (BW) were measured using 30cm fish measuring board and electronic scales.

Reproduction frequencies

Reproduction frequency of the Red claw Crayfish at Lake Kariba was determined using length frequency histograms constructed using Microsoft Excel, 2019 (Microsoft Corporation, 2019). The number of peaks on length-frequency Histograms were used to determine the reproduction frequency of sampled Crayfish.

Sex ratio

The sex ratio of Crayfish at Lake Kariba was determined by dividing the number of Females by the number of Males as earlier determined by Mukuka (2019).

Length-weight Relationships

The growth form coefficient (b) was determined by linear regression of the length-weight relationship (LWR) in Microsoft Excel, 2019 (Microsoft Corporation, 2019) using the equation: $W = aL^b$; where W = weight in grams, L = total length in centimeters, a is a scaling constant and b is the allometric growth form. This equation was logarithmically-transformed and expressed as: $\log \text{Weight} = a + b \log \text{Total length}$ (Froese, 2006). The growth exponent coefficient, r^2 , was used to denote the growth rate of the sampled specimens (Makeche *et al.*, 2023).

Growth performance indices

The Von Bertalanffy growth parameters k and L_∞ were estimated using the Ford-Walford, 1946 (Sparre and Venema, 1998) method and Pauly's equation of 1979 (Sparre and Venema, 1998), that is, $k = -\frac{1}{\Delta t} \times \ln b$ (Sparre and Venema, 1998); where b is a constant obtained by regression analysis of L(t) values of the sample size and Δt is change in time and $L_\infty = \frac{L_{\max}}{0.95}$ (Sparre and Venema, 1998); where L_{\max} is the maximum total length measurement recorded. Growth performance indices (Φ') were then estimated using the equation by Pauly and Munro (1984) expressed as: $\Phi' = \log_{10}(k) + 2 \log_{10}(L_\infty)$, where;

k = Von Bertalanffy growth coefficient and L_∞ = Von Bertalanffy asymptotic length.

Condition factors

Based on the expression of Fulton (1904): $K = 100 \times (W/L^3)$, where W is body weight (g), and L is Total Length (cm), Fulton's condition factor (K) was estimated. To obtain a Condition factor value close to unit, a scaling factor of 100 was used (Froese, 2006).

Natural mortality


The natural mortality (M_w) of *C. quadricarinatus* was determined from the equation: $M_w = 1.92 \text{ year}^{-1} \times (W)^{-0.25}$ (Peterson and Wroblewski (1984); where, M_w = natural mortality at mass W; and $W = a \times L^b$, a and b are the regression variables of length and weight (total length against body weight).

Statistical Analysis

The collected results were analyzed using linear regression and T-test using Microsoft Excel, 2019 (Microsoft Corporation, 2019). Linear regression was conducted using Microsoft Excel, 2019 (Microsoft Corporation, 2019) to determine the growth form coefficient 'b'. T-test analysis was conducted using Microsoft Excel, 2019 to determine significant differences ($P = 0.05$), if any, between the determined variates. One-way ANOVA was computed using Microsoft Excel, 2019 (Microsoft Corporation, 2019) to determine significant differences ($P = 0.05$), if any, among the regression variates.

Results

The display of the length-frequency histograms in Fig. 2 shows defined peaks, an indication that the *Cherax quadricarinatus* has well-defined spawning seasons in the Kafue Floodplain Fishery.






Fig. 2- Length-frequency histogram of *C. quadricarinatus* in the Kafue Floodplain Fishery.

The Histogram shows that *C. quadricarinatus* has a breeding frequency of 5 within one breeding season in the Kafue Floodplain Fishery.

Sex ratio

The sampled *C. quadricarinatus* species in the Kafue Floodplain Fishery were divided into 293 Females and 169 Males. This gave a sex ratio of 2 Females: 1 Male.

Male length-weight relationship and condition factor in the Kafue Floodplain Fishery

Male weight measurements ranged from 10.8g to 90.6g with a mean of 39g at in the Kafue Floodplain Fishery. Length measurements ranged from a low of 82mm to a high of 162mm with a mean of 121mm. The Fulton’s condition factor for Male *C. quadricarinatus* in the Kafue Floodplain Fishery was 2.2. There was significant correlation between Crayfish length and Crayfish weight (the adjusted r^2 was 0.864). ANOVA results showed a significant regression of length on weight ($P = 0.000$) (Table 1). The Linear regression equation of Crayfish length and Crayfish weight in the Kafue Floodplain Fishery was described by the mathematical model: $\log \text{Weight} = -4.29 + 2.81 \log \text{Total length}$. The growth form coefficient ‘b’ of Male *C. quadricarinatus* in the Kafue Floodplain Fishery was 2.81. The calculated ‘b’ value implies negative allometry because it is less than 3. The graphical representation of the length-weight relation is given in Fig. 3.

Table 1- Length-weight regression ANOVA results of Male *C. quadricarinatus* in the Kafue Floodplain Fishery

	<i>df</i>	<i>SS</i>	<i>MS</i>	<i>F</i>	<i>Significance F</i>
Regression	1	16.75108	16.75108	1489.809	3.7805E-132
Residual	372	4.182686	0.011244		
Total	373	20.93377			

df means degrees of freedom, *SS* means sum of squares, *MS* means mean sum of squares and *F* means the computer-generated comparison value. ANOVA means analysis of variance.




Fig. 3-Graph of length-weight relationship of Male *C. quadricarinatus* in the Kafue Floodplain Fishery.

Female length-weight relationship and condition factor in the Kafue Floodplain Fishery

Female weight measurements ranged from 12g to 98g with a mean of 34g in the Kafue Floodplain Fishery. Length measurements ranged from a low of 91mm to a high of 109mm with a mean of 120mm. The Fulton's condition factor for Female *C. quadricarinatus* in the Kafue Floodplain Fishery was 1.97. There was significant correlation between Female Crayfish length and Female Crayfish weight (adjusted $r^2 = 0.745$). ANOVA results showed a significant regression of length on weight ($P = 0.000$) (Table 2). The Linear regression equation of Female Crayfish length and Female Crayfish weight in the Kafue Floodplain Fishery was described by the mathematical model: $\log \text{Weight} = -4.01 + 2.66 \log \text{Total length}$. The allometric growth form 'b' of Female *C. quadricarinatus* in the Kafue Floodplain Fishery was 2.66. The calculated 'b' value implies negative allometry because it is less than 3. The graphical representation of the length-weight relation is given in Fig. 4.

Table 2-Length-weight regression ANOVA results of Female *C. quadricarinatus* in the Kafue Floodplain Fishery

	<i>df</i>	<i>SS</i>	<i>MS</i>	<i>F</i>	<i>Significance F</i>
Regression	1	4.682385	4.682385	848.919	2.84697E-88
Residual	291	1.60507	0.005516		
Total	292	6.287455			

df means degrees of freedom, *SS* means sum of squares, *MS* means mean sum of squares and *F* means the computer-generated comparison value. ANOVA means analysis of variance.




Fig. 4-Graph of length-weight relationship of Female *C. quadricarinatus* in the Kafue Floodplain Fishery.

Overall length-weight relationship and condition factor at Lake Kariba

Weight measurements ranged from 10.8g to 98g with a mean of 36g in the Kafue Floodplain Fishery. Length measurements ranged from a low of 82mm to a high of 162mm with a mean of 120mm. The Fulton's condition factor for combined data of *C. quadricarinatus* in the Kafue Floodplain Fishery was 2.20. There was significant correlation between Crayfish length and Crayfish weight (adjusted $r^2 = 0.780$). ANOVA results showed a significant regression of length on weight ($P = 0.000$) (Table 3). The Linear regression equation of Crayfish length and Crayfish weight in the Kafue Floodplain Fishery was described by the mathematical model: $\log \text{Weight} = -4.17 + 2.74 \log \text{Total length}$. The allometric growth form, 'b', of *C. quadricarinatus* in the Kafue Floodplain Fishery was 2.74. The calculated 'b' value implies negative allometry since it is less than 3. The graphical representation of the length-weight relation is given in Fig. 5.

Table 3-Length-weight regression ANOVA results of *C. quadricarinatus* in the Kafue Floodplain Fishery

	<i>df</i>	<i>SS</i>	<i>MS</i>	<i>F</i>	<i>Significance F</i>
Regression	1	8.265907	8.265907	1626.75	3.8042E-153
Residual	460	2.337371	0.005081		
Total	461	10.60328			

df means degrees of freedom, *SS* means sum of squares, *MS* means mean sum of squares and *F* means the computer-generated comparison value. ANOVA means analysis of variance.






Fig. 5-Graph of length-weight relationship of *C. quadricarinatus* in the Kafue Floodplain Fishery.

The Histograms which were constructed to show the Condition factors at the study sites within the Kafue Floodplain Fishery are shown in Fig. 6a and b.



(a)



(b)

Fig. 6- Histogram of Condition factors at Lake Kariba study sites (a) Sex Histogram, and (b) Study site.

Growth performance indices and natural mortality results

Growth performance indices ranged from a low of 4.3 among Female *C. quadricarinatus* at Kafue Road Bridge and Males at Chanyanya to a high 4.5 among Females at Chanyanya and among Males at Kafue Road Bridge (Table 5). All the growth performance indices were

supported by above-average growth rates (r^2 values above 0.5) at all study sites and across both gender. Natural mortality values of *C. quadricarinatus* ranged from a low of 0.0411 among Females at Chanyanya to a high of 0.0622 among Males at Kafue Road Bridge.

Table 5-Growth parameters, condition factors, growth performance indices and natural mortality parameters of *C. quadricarinatus* in the Kafue Floodplain Fishery.

	Kafue Road Bridge									Chanya								
	N	TL	W	a	b	r^2	K	M_w	Φ'	N	TL	W	a	b	r^2	K	M_w	Φ'
Male	161	121	39	-4.4	2.86	0.879	2.20	0.0622	4.5	8	120	33	-3.14	2.23	0.766	1.91	0.099	4.3
Female	274	120	34	-3.15	2.23	0.756	1.97	0.999	4.3	19	113	27	-4.617	2.93	0.929	1.87	0.0411	4.5
Total	435	121	36	-4.09	2.70	0.766	1.45	0.0530	4.9	27	126	29	-4.203	2.73	0.881	1.45	0.0141	4.4

N = number of *C. quadricarinatus* sampled, TL = mean total length in millimeters, W = mean weight in grammes, a = intercept, b = regression slope, r^2 = coefficient of determination, K = Fulton's condition factor, M_w = natural mortality and Φ' = growth performance index. Cha means Chanyanya study site.

All the growth coefficients (r^2) in the Kafue Floodplain Fishery were above 0.5. Condition factors at all study sites were all above-optimal ($K > 1$). The calculated value of the Condition factor shows that Crayfish is in good condition in the Kafue Floodplain Fishery. Natural mortality values were below-average at both study sites for combined data ($M_w < 0.5$). The value of the allometric growth format both study sites were less than 3, depicting negative allometry. Negative allometry implies that the increase in body weight did not match increase in total length

Discussion

The results of the current study on breeding frequency of *Cherax quadricarinatus* agree with the findings of Levi *et al.* (1997) who found that Red claw Crayfish breeding occurs naturally during the summer months, when temperatures are more than 20 °C. *C. quadricarinatus* mainly spawns during spring and summer while molting occurs mainly after the breeding season, but also between spawns (Karplus *et al.*, 1997; Levi *et al.*, 1997). Nunes *et al.* (2016) who reported that Crayfish normally breeds between September and April within their natural range in northern Australia freshwater bodies. The previous study by Makwelele (2017) in the Kafue Floodplain established that *C. quadricarinatus* has well-defined spawning seasons.

The sex ratio results are similar to the results obtained in the Upper Zambezi by Mukuka (2019), who also found a sex ratio of almost one to one. Both studies established that there are more Males than Females in most aquatic water bodies. The sex ratio results are also in agreement with those found in Mexico by Bortolini *et al.* (2007) and Belle *et al.* (2011) in Singapore. However, they are different from those found in Swaziland, South Africa, and Zimbabwe (Nunes *et al.*, 2017b; Marufu *et al.*, 2018) where Females were outnumbering Males.

The growth exponent coefficient (r^2) of *C. quadricarinatus* was above-average (more 0.5) at both study sites, implying maximum growth rate. These results are characteristic of invasive species as earlier found in *Oreochromis niloticus* by Chikopela *et al.* (2011), Makeche *et al.* (2023) and Nyirenda *et al.* (2024) in Zambia. The large growth exponents were supported by large growth performance indices at both study sites. These results are in agreement with earlier results by Marufu *et al.* (2018) ($\Phi' = 3.96$) from Lake Kariba, Mukuka (2019) ($\Phi' = 4.0$) from the Upper Zambezi and Nyirenda *et al.* (2024) who found similar values among *Oreochromis niloticus*, an invasive fish at Lake Kariba ($\Phi' = 4.58$). Makwelele (2017) found lower results ($\Phi' = 2.8$) from another part of Zambia. The difference between the current results and those obtained by Makwelele (2017) can be attributed to both the habitat and the sampling period. The current study was conducted during the rainy season when food materials are abundant while Makwelele (2017) conducted his study during the dry season when most natural habitats are nutrient-poor. The sampling season may also explain why Condition factor values of the present study were larger than those obtained by Makwelele (2017) ($K = 1.8$ to 2.3). These above-average values show that *C. quadricarinatus* is well acclimatized to its habitat (Makeche *et al.* (2023). The value of the Condition factor is a good indicator of the productivity of the habitat, ecological conditions and biological factors such as fatness, gonad maturity and habitat adaptability (Saha *et al.*, 2021).

The Natural mortality rate estimates obtained in this study are similar to those obtained by Makwele (2017) ($M = 0.06$ to 0.08) from the Kafue Floodplains and Makeche *et al.* (2020) on another invasive organism. The results of the current study are however different and lower than those those obtained by Marufu *et al.* (2018) ($M = 0.99$) from Lake Kariba and those found by Mukuka (2019). This may be due to several reasons. Firstly, it is always difficult to capture a representative length-frequency sample (real size distribution of the population) even with a large sample size due to gear selectivity. This has a compounding effect on the Von Bertalanfy growth equation parameter estimations which Mukuka (2019) used in estimating the growth coefficient, natural mortality, total mortality and asymptotic length. Also, the length-converted catch curve method which Mukuka (2019) used tends to over-estimate the total mortality when seasonal growth is considered (Gayanilo and Pauly, 1997).

Conclusion

The success of *C. quadricarinatus* in invading and establishing itself on Lake Kariba may not absolutely reflect on the species' inherently strong life-history strategies, instead, a number of external human-driven influences may have aided its current and future spread Lake Kariba. *C. quadricarinatus* is well-adapted to its habitat at Lake Kariba, as the growth exponents and condition factor results show. Although fish farmers may consider Crayfish to be a nuisance, it can be cultivated on a large scale as a reliable source of proteins while improving the economic returns of aquaculture farmers.

Acknowledgements

This study was sponsored by USAID.

REFERENCES :

1. Amponsah, K.K.S. Asiedu, B. and Failler, P. 2020. Population parameters of *Oreochromis niloticus* (L) from a semi-open lagoon (Sakumo II), Ghana and its implications on management. *Egyptian Journal of Aquatic Biology and Fisheries*, 24(2):195-207.
2. Austin, C.M., Jones, C. and Wingfield, M. 2010. *Cherax quadricarinatus*, Tropical Blue Crayfish. *The IUCN Red List of Threatened Species 2010*, no. e.T4621A11041003: 9.
3. Belle, C.C., Tan, H.H., Tan, S.H., Yeo, D.C.J., Todd, P.A., Clews, E. and Wong, J.Q.H. 2011. Ornamental Trade as a Pathway for Australian Red claw Crayfish Introduction and Establishment. *Aquatic Biology*, 12(1): 69–79.
4. Bortolini, R., José, L., Fernando, A. and Gabino, R-A. 2007. On the Presence of the Australian Red claw Crayfish, *Cherax quadricarinatus*, in Mexico. *Biological Invasions*, 9(5): 615–20.
5. Chikopela, S.T., Katongo, C. and Mudenda, H.G. 2011. Abundance of Mouth Brooding Tilapiines in the Kafue Floodplains, Zambia. *Journal of Ecology and the Natural Environment*, 3(10): 344-350.
6. Cowx, I.G. and Kalonga, M. 2018. Optimising hydropower development and ecosystem services in the Kafue River, Zambia. *Marine and Fresh water Research*, 69 (12): 1974-1982.
7. FAO. 2011. 'Cultured Aquatic Species Information Programme: *Cherax quadricarinatus* (von Martens, 1868)'. FAO Fisheries and Aquaculture Department, Rome, Italy.
8. Froese, R. 2006. Cube law, Condition factor and weight length-relationship: history meta-analysis and recommendations. *Journal of Applied Ichthyology*, 22(4): 241-253.
9. Fulton, T.W. 1904. The Rate of Growth of Fishes. Twenty-second Annual Reports, Part III. Fisheries Board of Scotland, Edinburgh pp. 141-241.
10. Gayanilo, F.C. and Pauly, D. 1997. FAO-ICLARM Stock Assessment Tools Reference Manual 39. *Stock Assessment Tools Reference Manual*, 39: 262-265.
11. Jaklič, M. and Vrezec, A. 2011. The First Tropical Alien Crayfish Species in European Waters: The Red claw *Cherax quadricarinatus* (Von Martens, 1868) (Decapoda, Parastacidae). *Crustaceana*, 84(6): 651–665.
12. Karplus, I., Michael, P.M., Barki, A. an, David, B.R. 1997. Australian Red Claw Crayfish, *April 1997*, SRAC Publication No. 244.
13. Levi, T., Haluta, G., Barki, A. and Karplus, I. 1997. Annual cycle of spawning and molting in the Red claw Crayfish, *Cherax quadricarinatus*, under laboratory conditions. *Aquaculture volume*, 157 (4): 139-149.
14. Makeche, M.C., Mudenda, H.G. and Katongo, C. 2020. Growth and Mortality of Mouth Brooding Tilapiines of the Kafue Floodplain Fishery, Zambia. *American Scientific Research Journal for Engineering, Technology, and Sciences*, 64: 109-118.
15. Makeche, M.C., Nhwatiwa, T., Chitondo, L., Kanyati, M., Katongo, C., Kamins,a C., Kaona, M., Ndebe, N., Mulavu, M., Khumalo, C.S, Simulundu, E., Changula, K., Chitanga, S., Mubemba, B., Muleya, W and, Makwelele, G. 2023. Comparative study of growth rates, condition factors and natural mortality of *Oreochromis niloticus* fish from culture fisheries and capture fisheries at Lake Kariba, Zambia. *International Journal of Fisheries and Aquaculture*, 15(2): 21-35.
16. Makwelele, G. 2017. Reproductive biology, growth and mortality of the introduced red claw crayfish, *Cherax quadricarinatus* in Kafue Floodplain Fishery. (M.Sc. dissertation). University of Zambia, Lusaka, Zambia.
17. Marufu, L., Barson, M., Chifamba, P., Tiki, M. and Nhwatiwa, T. 2018. The Population Dynamics of a Recently Introduced Crayfish, *Cherax quadricarinatus* (von Martens, 1868), in the Sanyati Basin of Lake Kariba, Zimbabwe. *African Zoology*, 53(1): 17–22.
18. Microsoft Corporation. 2019. Microsoft Excel User's manual. Analytical Software. Tallahassee, USA.

19. Mukuka, K.N. 2019. Establishment and Population Dynamics of the Alien Invasive Red Claw Crayfish (*Cherax quadricarinatus* Von Martens, 1868) on the Upper Zambezi River, Zambia. (M.Sc. dissertation). University of Bremen, Bremen, Germany.
20. Nunes, A.L., Robert, R.J., Ben, T., Measey, G.J., Olaf, L.F. and Weyl, G. 2016. Invasive Crayfish Threaten Okavango Delta. *Frontiers in Ecology and the Environment*, 14 (5): 237–38.
21. Nunes, A.L., Andries, C., Hoffman, A.C., Olaf, L.F., Weyl, G., Measey, G.J. and Zengeya, T.A. 2017b. Distribution and Establishment of the Alien Australian Red claw Crayfish, *Cherax quadricarinatus*, in South Africa and Swaziland. *PeerJ*, 5: e3135.
22. Nyirenda, M.S., Katongo, C., Mudenda, H.G., Mbewe, M. and Sinyinza, D. 2024. Aspects of Growth and Reproductive Biology among three Tilapiines on Lake Kariba. *International Journal of Research Publication and Reviews*, 5(1): 4294-4299.
23. Osho, E.F. and Usman, A.R. 2019. Length-weight relationship, condition factor and fecundity of African Snakehead *Parachanna obscura* from the Anambra River, South-East Nigeria. *Croatian Journal of Fisheries*, 77(2): 99-105.
24. Pauly, D. and Munro, J.L. 1984. Once more on Growth Comparison in Fish and Invertebrates. *Fishbyte*, 2(1): 21.
25. Peterson, I. and Wroblewski, J.S. 1984. Mortality rates of Fishes in the pelagic ecosystem. *Canadian Journal of Fisheries and Aquatic Sciences*, 41:1117-1120.
26. Ramsar. 2007. The Annotated List: Zambia, Kafue Flats. Available at <https://rsis Ramsar.org/ris/530>.
27. Saha, N., Roy, P., Nadia, M.Z., Ahamed, F. and Hossain, Y.M. 2021. Growth, Condition, Maturity and Mortality of the Dwarf Gourami, *Trichogaster lalius* (Hamilton, 1822) in a Wetland Ecosystem (Beel Dakatia), Southwestern Bangladesh. *Egyptian Journal of Aquatic Biology and Fisheries*, 25(3): 505-524.
28. Sparre, P. and Venema, S.C. 1998. Introduction to Tropical Fish Stock Assessment, Part: 1 Manual, *FAO Fisheries Technical Paper 306/1, Rev 2*, Rome.
29. Tyser, A.B. and Douthwaite, J.B. 2014. Predation on invasive Red claw crayfish *Cherax quadricarinatus* by native fishes in the Kafue River, Zambia. *African Journal of Aquatic Sciences*, 39 (4): 473-477